Bone Grafting and Barrier Membrane Consent Form

Procedure Planned: I understand that bone gaffing and barrier membrane procedures include inherent risks such as but not limited to the following:	
	Infection . No matter how carefully surgical sterility is maintained, it is possible, because of the existing non-sterile oral environment, for infections to occur postoperatively. At times, these may be of a serious nature. Should severe swelling occur, particularly accompanied with fever or malaise, professional attention should be received as soon as possible.
	Bleeding, bruising, and swelling . Some moderate bleeding may last several hours. If profuse, you must contact us as soon as possible. Some swelling is normal, but if severe, you should notify us. Swelling usually starts to subside after about 48 hours. Bruises may persist for a week or so.
	Loss of all or part of the graft. Success with bone and membrane grafting is high. Nevertheless, it is possible that the graft could fail. A block bone graft taken from somewhere else in your mouth may not adhere or could become infected. Despite meticulous surgery, particulate bone graft material can migrate out of the surgery site and be lost. A membrane graft could start to dislodge, If so, the doctor should he notified. Your compliance is essential to assure success.
5.	Types of graft material . Some bone graft and membrane material commonly used are derived from human or other mammal sources. These grafts are thoroughly purified by different means to be free from contaminants. Signing this consent form gives your approval for the doctor to use such materials according to his knowledge and clinical judgment for your situation.
7.	Injury to nerves . This would include injuries causing numbness of the lips; the tongue; any tissues of the mouth; and/or cheeks or face. This numbness which could occur, may be of a temporary nature, lasting a few days, a few weeks, a few months, or could possibly be permanent, and could be the result of surgical procedures or anesthetic administration. Sinus involvement . In some cases, the root tips of upper teeth lie close in proximity to the maxillary sinus. Occasionally, with extractions and/or grafting near the sinus, the sinus can become involved. If this happens, you will need to take special medications. Should sinus penetration occur, it may be necessary to later have the sinus surgically closed.
	responsibility to seek attention should any undue circumstances occur post-operatively and you should diligently follow operative and post-operative instructions.
Informed Consent : As a patient, I have been given the opportunity to ask any questions regarding the nature and purpose of surgical treatment and have received answers to my satisfaction. I do voluntarily assume any and all possible risks, including the risk of harm, if any, which may be associated with any phase of this treatment in hopes of obtaining the desired results, which may or may not be achieved. No guarantees or promises have been made to me concerning my recovery and results of the treatment to be rendered to me. The fee(s) for this service have been explained to me and are satisfactory. By signing this form, I am freely giving my consent to allow and authorize Dr. Joiner, Dr. Zwart and associates to render any treatment necessary or advisable to my dental conditions, including any and all anesthetics and/or medications.	

Witness signature

Date

Patient (Parent) signature

Date